

Table 1. Estimated HIV incidence among persons aged ≥13 years, by year of infection and selected characteristics, 2018–2022—United States (cont)

	No.	RSE (%)	95% CI	Rate ^a	95% CI
2020 (COVID-19 pandemic)^j					
Sex at birth					
Male	28,400	3.5	26,400–30,300	20.5	19.2–21.9
Female	5,800	6.9	5,000–6,600	4.1	3.6–4.7
Age at infection (yr)					
13–24	7,600	6.6	6,600–8,500	14.4	12.5–16.2
25–34	13,700	4.9	12,400–15,000	30.0	27.2–32.9
35–44	6,600	7.0	5,700–7,600	15.5	13.4–17.7
45–54	3,700	9.4	3,000–4,400	9.0	7.3–10.6
55–64	2,100	12.2	1,600–2,600	5.0	3.8–6.2
≥65	480	25.9	240–720	0.9	0.4–1.3
Race/ethnicity					
American Indian/Alaska Native	230	*36.3	70–390	11.5	3.3–19.7
Asian	480	26.4	230–720	2.8	1.4–4.3
Black/African American	13,600	4.7	12,300–14,800	39.3	35.7–42.9
Hispanic/Latino ^b	10,000	6.5	8,700–11,300	20.7	18.0–23.3
Native Hawaiian/other Pacific Islander
White	8,600	5.6	7,600–9,500	5.0	4.4–5.5
Multiracial	1,200	16.4	830–1,600	24.0	16.3–31.7
Transmission category^c					
Male-to-male sexual contact ^d	23,200	3.8	21,400–24,900	—	—
Injection drug use ^e	2,600	10.5	2,100–3,200	—	—
Male	1,700	14.0	1,200–2,100	—	—
Female	970	15.2	680–1,300	—	—
Male-to-male sexual contact ^d and injection drug use ^e	1,400	12.9	1,000–1,700	—	—
Heterosexual contact ^f	6,900	7.1	6,000–7,900	—	—
Male	2,100	15.4	1,500–2,700	—	—
Female	4,800	7.7	4,100–5,600	—	—
Region of residence^g					
Northeast	4,500	8.7	3,700–5,300	9.2	7.6–10.7
Midwest	4,800	7.9	4,100–5,600	8.3	7.0–9.6
South	17,100	4.4	15,600–18,600	16.1	14.7–17.5
West	7,700	6.7	6,700–8,800	11.7	10.2–13.2
Total^h	34,200	3.1	32,100–36,300	12.2	11.5–13.0
2021ⁱ					
Sex at birth					
Male	26,500	4.0	24,400–28,600	19.1	17.6–20.6
Female	6,200	7.1	5,300–7,000	4.3	3.7–4.9
Age at infection (yr)					
13–24	6,800	7.7	5,800–7,800	12.8	10.9–14.8
25–34	13,000	5.6	11,600–14,400	28.6	25.5–31.7
35–44	6,700	7.7	5,700–7,700	15.4	13.1–17.7
45–54	3,600	10.5	2,800–4,300	8.8	7.0–10.6
55–64	2,100	13.4	1,600–2,700	5.0	3.7–6.3
≥65	540	27.7	250–830	1.0	0.4–1.5
Race/ethnicity					
American Indian/Alaska Native	180	*46.4	20–340	8.7	0.8–16.7
Asian	430	*31.1	170–690	2.5	1.0–4.1
Black/African American	13,000	5.3	11,700–14,400	37.4	33.6–41.3
Hispanic/Latino ^b	9,800	7.2	8,400–11,200	19.9	17.1–22.8
Native Hawaiian/other Pacific Islander
White	8,100	6.3	7,100–9,100	4.7	4.1–5.3
Multiracial	1,100	19.2	660–1,500	20.1	12.5–27.6
Transmission category^c					
Male-to-male sexual contact ^d	21,600	4.4	19,700–23,500	—	—
Injection drug use ^e	2,600	12.1	1,900–3,200	—	—
Male	1,500	17.1	1,000–2,000	—	—
Female	1,100	16.5	710–1,400	—	—
Male-to-male sexual contact ^d and injection drug use ^e	1,300	14.5	920–1,600	—	—
Heterosexual contact ^f	7,200	7.4	6,100–8,200	—	—
Male	2,100	16.5	1,400–2,800	—	—
Female	5,100	7.9	4,300–5,900	—	—
Region of residence^g					
Northeast	4,100	10.0	3,300–4,900	8.4	6.7–10.0
Midwest	4,700	8.9	3,900–5,500	8.1	6.7–9.5
South	16,700	4.9	15,100–18,300	15.6	14.1–17.1
West	7,100	7.7	6,100–8,200	10.7	9.1–12.4
Total^h	32,700	3.5	30,400–34,900	11.6	10.8–12.4